UNITED STATES DEPARTMENT OF AGRICULTURE NATURAL RESOURCES CONSERVATION SERVICE ECOLOGICAL SITE DESCRIPTION

ECOLOGICAL SITE CHARACTERISTICS

Site Type: Rangeland	
Site ID: R070XC104NM	
Site Name: Deep Sand	
Precipitation or Climate Zone:	13 to 16 inches
Phase:	

PHYSIOGRAPHIC FEATURES

٦	×Τ				4.		
ı	N	a	r	ra	Ш	\mathbf{v}	e

This site occurs as the coarse-textured eolian and alluvial sediments on the upland plains. Slopes are nearly level to gently undulating, generally less than 5 percent. Low stabilized hummocks or dunes frequently occur. Exposure varies and is not significant. Elevations range from 5,000 to 7,000 feet above sea level.

|--|

- 1. Plain
- 2. Sand sheet
- 3.

Aspect:

- 1. N/A
- 2.
- 3.

	Minimum	Maximum
Elevation (feet)	5,000	7,000
Slope (percent)	0	5
Water Table Depth (inches)	N/A	N/A

Flooding:MinimumMaximumFrequencyN/AN/ADurationN/AN/A

Ponding:MinimumMaximumDepth (inches)N/AN/AFrequencyN/AN/ADurationN/AN/A

Runoff Class:

Negligible to medium.

CLIMATIC FEATURES

Narrative:

The average annual precipitation ranges from 13 to 16 inches. Variations of 5 inches, more or less, are not uncommon. Seventy-five percent of the precipitation falls from April to October. Most of the summer precipitation comes in the form of high intensity-short duration thunderstorms.

Temperatures are characterized by distinct seasonal changes and large annual and diurnal temperature changes. The average annual temperature is about 50 degrees F with extremes of -29 in the winter and 103 degrees F in the summer.

The average frost-free season is 130 to 160 days. The last killing frost is in early May and the first killing frost is in early October.

Both temperature and precipitation favor warm-season species, However, about 40 percent of the precipitation falls at a time favorable to cool-season plant growth. This allows the cool-season plants to occupy an important component of this site. Because of the soils on this site, the plant community can respond quickly to any precipitation during the frost-free period. Also, because of the soil, strong winds that blow from the west and southwest from February to June can dry the soil profile rapidly during a critical period for plant growth. These strong winds can also cause the soil to blow. Wind-blown particles cause plant damage and reduce growth.

Climate data was obtained from http://www.wrcc.sage.dri.edu/summary/climsmnm.html web site using 50% probability for freeze-free and frost-free seasons using 28.5 degrees F and 32.5 degrees F respectively.

	Minimum	Maximum
Frost-free period (days):	131	173
Freeze-free period (days):	155	187
Mean annual precipitation (inches):	13	16

Monthly moisture (inches) and temperature (⁰F) distribution:

J	Precip. Min.	Precip. Max.	Temp. Min.	Temp. Max.
January	.34	.92	15.6	42.1
February	.34	.81	19.9	52.9
March	.23	.98	24.4	59.7
April	.39	.96	31.4	68.9
May	.85	1.61	39.2	77.7
June	.89	1.62	46.9	87.1
July	1.77	2.75	53.1	88.5
August	2.46	3.22	51.9	85.7
September	1.54	2.26	44.3	80.4
October	1.0	1.51	32.8	70.5
November	.57	1.02	22.2	57.5
December	.34	1.16	15.9	49.3

Climate Sta	ntions:						
					Period	f	
Station ID	291918	Location	Clines Corners7SE, NM	From:	12/10/68	To:	11/30/00
Station ID	292096	Location	Coronal 1SSW, NM	From:	12/1/77	To:	9/30/92
Station ID	293060	Location	Estancia, NM	From:	01/01/14	To:	12/31/00
Station ID	293649	Location	Gran Quivira Natl. Monument, NM	From:	06/01/38	To:	12/31/00
Station ID	295965	Location	Mountainair, NM	From:	03/01/14	To:	12/31/00
Station ID	299405	Location	Vaughn, NM	From:	01/01/71	To:	12/31/00

INFLUENCING WATER FEATURES

Narrative:

This site is not influenced by water from wetlands or streams.

Wetland description:

System	Subsystem	Class
N/A		

If Riverine Wetland System enter Rosgen Stream Type:
N/A

REPRESENTATIVE SOIL FEATURES

Narrative:

The soils on this site are deep and excessively drained. The surface textures are loamy fine sand and sand that extend to a depth of 60 inches or more. The soils are rapidly permeable and a low water-holding capacity. Surface runoff is very slow. Drying surface is fast and soil blowing hazard is high.

Parent Material Kind: Eolian Sands

Parent Material Origin: Sandstone - unspecified

Surface Texture:

Loamy fine sand
 Sand
 .

Surface Texture Modifier:

1. N/A	
2.	
3.	

Subsurface Texture Group: Loamy fine sand

Surface Fragments <=3" (% Cover): N/A

Surface Fragments >3" (% Cover): N/A

Subsurface Fragments <= 3" (%Volume): N/A
Subsurface Fragments <= 3" (%Volume): N/A

	Minimum	Maximum
Drainage Class:	Excessively	
Permeability Class:	Rapid	+
Depth (inches):	60	>72
Electrical Conductivity (mmhos/cm):	0.00	2.00
Sodium Absorption Ratio:	N/A	N/A
Soil Reaction (1:1 Water):	7.4	8.4
Soil Reaction (0.1M CaCl2):	N/A	N/A
Available Water Capacity (inches):	3	6
Calcium Carbonate Equivalent (percent):	N/A	N/A

PLANT COMMUNITIES

Ecological Dynamics of the Site :	
Beological Bynamics of the Site.	
Plant Communities and Transitional Pathways (di	noram)
Zame Communication and Limited Mark and Marks (Miles	

Plant Community Nan	ne: Historic Climax Pl	ant Community						
Plant Community Seq	uence Number: 1	Narrative Label:	НСРС					
Plant Community Narrative: Historic Climax Plant Community The site is a grassland characterized by both warm- and cool-season perennial tall and midgrasses, low growing shrubs and half-shrubs, and a variety of forbs. Forb poduction fluctuates greatly from year to year. In years of significant spring and fall moisture, forb composition and production are a very important part of the plant community.								
Canopy Cover:								
Trees		$\frac{0-1 \%}{5-8 \%}$	_					
Shrubs and Half-shrubs			5 – 8 %					
	e Percent of Surface Area							
Grasses & Forbs		$ \begin{array}{r} 15 - 20 \\ \hline 55 - 65 \\ \hline 0 - 1 \end{array} $	15 – 20					
Bare ground		55 – 65	55 – 65					
Surface cobble and ston	e	0 - 1	0 - 1					
Litter (percent)		8 – 10						
Litter (average depth in	cm.)	_ 2						
Plant Community Ann	nual Production (by plan	nt type):						
	Annual Produ	iction (lbs/ac)						
Plant Type	Low	RV	High					
Grass/Grasslike	200	940	1,680					
Forb	20	94	168					
Tree/Shrub/Vine	20	94	168					
Lichen								

1,175

250

Moss

Totals

Microbiotic Crusts

2,100

Plant Community Composition and Group Annual Production:

Plant Type - Grass/Grasslike

Group	Scientific	assinc	Species Annual	Group Annual
Number	Plant Symbol	Common Name	Production Production	Production
1	BOGR2	Blue Grama	118 - 176	118 – 176
	PLJA	Galleta		
2	BOER4	Black Grama	59 - 118	59 – 118
3	SCSC	Little Bluestem	118 - 294	118 - 294
	ANGE	Big Bluestem		
	BOBA3	Cane Bluestem		
	ANHA	Sand Bluestem		
4	ACHY	Indian Ricegrass	118 - 176	118 - 176
5	BOCU	Sideoats Grama	59 - 118	59 – 118
6	PAVI2	Switchgrass	59 - 118	59 – 118
	SONU2	Indiangrass		
	CAGI3	Giant Sandreed		
7	HENE5	New Mexico Feathergrass	35 - 59	35 - 59
	HECO26	Needleandthread		
8	SPCR	Sand Dropseed	118 -176	118 - 176
	SPFL2	Mesa Dropseed		
	SPCO4	Spike Dropseed		
	SPGI	Giant Dropseed		
9	2GRAM	Other Grasses	59 - 118	59 - 118

Plant Type - Forb

I mile I J P				
Group	Scientific		Species Annual	Group Annual
Number	Plant Symbol	Common Name	Production	Production
10	ERIOG	Wildbuckwheat	12 - 35	12 - 35
11	HEAN3	Annual Sunflower	12 - 35	12 - 35
12	CROTO	Croton	12 - 35	12 - 35
13	2FORBS	Other Forbs	12 - 35	12 - 35

Plant Type – Tree/Shrub/Vine

Tiant Typ	e = Tree/Sin	ID/ VIIIC		
Group	Scientific		Species Annual	Group Annual
Number	Plant Symbol	Common Name	Production	Production
14	ARFI2	Sand Sagebrush	59 -118	59 – 118
	ARBI3	Bigelow Sagebrush		
15	RHTR	Skunkbush Sumac	24 - 59	24 - 59
	RHMI3	Littleleaf Sumac		
16	DAFO	Feather Dalea	12 - 35	12 - 35
17	2SD	Other Shrubs	12 - 35	12 - 35

Plant Type - Lichen

•	Group Number	Scientific Plant Symbol	Common Name	Species Annual Production	Group Annual Production

Plant Type - Moss

Group Number	Scientific Plant Symbol	Common Name	Species Annual Production	Group Annual Production

Plant Type - Microbiotic Crusts

Group	Scientific		Species Annual	Group Annual
Number	Plant Symbol	Common Name	Production	Production

Other grasses that could appear on this site include: alkali sacaton, threeawn spp., sandhill muhly, purple lovegrass, ring muhly, bottlebrush squirreltail, western wheatgrass, plains bristlegrass, green sprangeltop, and bush muhly.

Other shrubs include: ephedra spp., winterfat, rabbitbrush, broom snakeweed, fourwing saltbush, yucca spp., cacti spp., juniper, pinyon, algerita, and oak spp..

Other forbs include: Tansymustard, locoweed spp., red stem milkvetch, scarlet globemallow, mariola, sand verbena, goldenweed, and threadleaf groundsel.

Plant Growth Curves

Growth Curve ID 4304NM

Growth Curve Name: **HCPC**

Growth Curve Description: Mixed warm and cool-season mid and tall perennial grassland

with shrubs and half-shrubs and forbs.

Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
0	0	5	7	10	15	25	25	8	5	0	0

ECOLOGICAL SITE INTERPRETATIONS

Animal Community:

Habitat for Wildlife:

This range site produces a habitat which supports a resident animal community characterized by pronghorn antelope, kit fox, badger, desert cottontail, spotted ground squirrel, southern plains woodrat, Ord's kangaroo rat, plains pocket mouse, northern grasshopper mouse, Botta's pocket gopher, sparrow hawk, mourning dove, meadowlark, chipping sparrow, plains spadefoot toad, eastern fence lizard, plateau whiptail, short-horned lizard, and prairie rattlesnake. Common raven and prairie falcon hunt over this site. Where dense stands of large pinyon, juniper or ponderosa pine occur, woodland wildlife species such as mule deer, gray fox, rock squirrel, Harlequin quail, pinyon jay, scrub jay, and Cassin's kingbird become characteristic.

Hydrology Functions:

The runoff curve numbers are determined by field investigations using hydrologic cover conditions and hydrologic soil groups.

Herdusla eia Intampatationa							
Hydrologic Interpretations							
Soil Series	Hydrologic Group						
Benevides	A						
Mido	A						
Valent	A						

Recreational Uses:

This site offers fair potential for hiking, horseback riding, nature observation, photography, camping and picnicking. It also offers fair to good opportunity for hunting pronghorn antelope, mourning dove, quail, and varmints. Trapping fur-bearing animals is good. Natural beauty is dependent upon scattered flowering shrubs and forbs and the open grassland character of the site.

Wood Products:

This site has no significant potential for wood products.

Other Products:

Grazing:

This site is suitable for grazing by all kinds and classes of livestock during all seasons of the year. This site can respond rapidly to good management and responds best to a system of grazing which rotates the season of use. This site is poorly suited for continuous year-long grazing. If grazing is continued year long or grazing pressure is too heavy, the dominant grasses such as black grama, little bluestem, cane bluestem, sand bluestem, Indian ricegrass, and sideoats grama will decrease rapidly. This will cause a corresponding increase in woody species, annuals, and grasses like threeawn, sand dropseed, sandhill muhly, and ring muhly. This will also cause a decrease in the forage production. Severe site deterioration is characterized by increased amounts of bare soil which causes soil blowing and hummocking. The site is sometimes invaded by woody species such as pinyon and juniper, or in rare cases, ponderosa pine, and may support relatively long-lived stands of these species.

Other Information:							
Guide to Suggested Initial Stocking Rate Acres per Animal Unit Month							
Similarity Index	Ac/AUM						
100 - 76	2.0 - 2.8						
75 – 51	2.6 - 4.4						
50 – 26	3.3 - 7.0						
25 – 0	7.0+						

Plant Part	Code	Species Preference	Code
Stems	S	None Selected	NS
Leaves	${f L}$	Preferred	P
Flowers	F	Desirable	D
Fruits/Seeds	F/S	Undesirable	U
Entire Plant	EP	Not Consumed	NC
Underground Parts	UP	Emergency	E
		Toxic	T

Plant Preference by Animal Kind:

Animal Kind: Livestock
Animal Type: Cattle

		Plant		Forage Preferences										
Common Name	Scientific Name	Part	J	F	M	A	M	J	J	A	S	О	N	D
Black Grama	Bouteloua eriopoda	EP	P	P	P	D	D	D	D	D	D	D	P	P
Little Bluestem	Schizachyrium scoparium	EP	D	D	D	P	P	P	P	D	D	D	D	D
Sand Bluestem	Andropogon hallii	EP	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S
Indian Ricegrass	Achnatherum hymenoides	EP	P	P	P	P	P	P	P	P	P	P	P	P
Sideoats Grama	Bouteloua curtipendula	EP	P	P	P	P	P	P	P	P	P	P	P	P
Big Bluestem	Andropogon gerardii	EP	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S
Switchgrass	Panicum virgatum	EP	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S
New Mexico Feathergrass	Hesperostipa neomexicana	EP	D	D	P	P	P	D	D	D	D	D	D	D
Needleandthread	Hesperostipa comata	EP	D	D	P	P	P	D	D	D	D	D	D	D
Annual Sunflower	Helianthus annuum	EP	U	U	U	U	U	D	D	D	U	U	U	U

Animal Kind: Livestock
Animal Type: Sheep

		Plant	Forage Preferences											
Common Name	Scientific Name	Part	J	F	M	A	M	J	J	A	S	О	N	D
Black Grama	Bouteloua eriopoda	EP	D	D	D	D	D	D	D	D	D	D	D	D
Indian Ricegrass	Achnatherum hymenoides	EP	P	P	D	D	D	D	D	D	D	P	P	P
Sideoats Grama	Bouteloua curtipendula	EP	D	D	D	D	D	D	D	D	D	D	D	D
Bigelow Sagebrush	Artemisia bigelovii	L/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S
Annual Sunflower	Helianthus annuum	EP	U	U	U	U	U	D	D	D	U	U	U	U

Animal Kind: Wildlife
Animal Type: Antelope

		Plant	Forage Preferences											
Common Name	Scientific Name	Part	J	F	M	A	M	J	J	A	S	О	N	D
Indian Ricegrass	Achnatherum hymenoides	EP	U	U	P	P	P	U	U	U	P	P	P	U
Wildbuckwheat	Eriogonum species	EP	U	U	D	D	D	D	D	D	U	U	U	U
Annual Sunflower	Helianthus annuum	EP	U	U	U	U	U	D	D	D	U	U	U	U
Skunkbush Sumac	Rhus trilobata	L/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S

SUPPORTING INFORMATION

Associated sites: Site Name Site ID Site Narrative Similar sites: Site Name Site ID Site Narrative **State Correlation**: This site has been correlated with the following sites: **Inventory Data References: Data Source** # of Records Sample Period County State **Type Locality**: **State:** New Mexico County: Chavez, De Baca, Guadalupe, Lincoln, San Miguel, Santa Fe, Torrance Latitude: Longitude: Township: Range: Section: Is the type locality sensitive? No Yes **General Legal Description**: **Relationship to Other Established Classifications**: **Other References:** Data collection for this site was done in conjunction with the progressive soil surveys within the Pecos-Canadian Plains and Valleys 70 Major Land Resource Area of New Mexico. This site has been mapped and correlated with soils in the following soil surveys: Chaves, De Baca, Guadalupe, Lincoln, Sna Miguel, Santa Fe, Torrance. **Characteristic Soils Are:** Valent Benevides Other Soils included are: Mido **Site Description Approval:** Author Date Approval Donald H. Fulton 03/03/82 Don Sylvester 02/02/82 **Site Description Revision:** Author Approval Date Date Elizabeth Wright 12/17/02 06/19/01 George Chavez